

A Boundedness Result for Minimizers of Some Polyconvex Integrals

Menita Carozza, Hongya Gao, Raffaella Giova & Francesco Leonetti

Journal of Optimization Theory and Applications

ISSN 0022-3239

J Optim Theory Appl

DOI 10.1007/s10957-018-1335-0

Vol. 177, No. 3

**ONLINE
FIRST**

JOURNAL OF OPTIMIZATION THEORY AND APPLICATIONS

Special Issue on
Recent Advances in Vector Optimization and Image Space Analysis
in Celebration of Prof. H.P. Benson's 68th Birthday

Guest Editors:

Guangya Chen, Chinese Academy of Sciences, China

Shengjie Li, Chongqing University, China

Jiuping Xu, Sichuan University, China

Xinmin Yang, Chongqing Normal University, China

 Springer

 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media, LLC, part of Springer Nature. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

A Boundedness Result for Minimizers of Some Polyconvex Integrals

Menita Carozza¹ · Hongya Gao² · Raffaella Giova³ · Francesco Leonetti⁴ 

Received: 21 March 2018 / Accepted: 8 June 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract We consider polyconvex functionals of the Calculus of Variations defined on maps from the three-dimensional Euclidean space into itself. Counterexamples show that minimizers need not to be bounded. We find conditions on the structure of the functional, which force minimizers to be locally bounded.

Keywords Local · Bounded · Minimizer · Polyconvex · Integral

Mathematics Subject Classification 49N60 · 35J50

Communicated by Bernard Dacorogna.

✉ Francesco Leonetti
leonetti@univaq.it

Menita Carozza
carozza@unisannio.it

Hongya Gao
578232915@qq.com

Raffaella Giova
raffaella.giova@uniparthenope.it

¹ Università del Sannio, Benevento, Italy

² Hebei University, Baoding, China

³ Università di Napoli “Parthenope”, Naples, Italy

⁴ Università di L’Aquila, L’Aquila, Italy

1 Introduction

Let us consider polyconvex integrals of the Calculus of Variations. Partial regularity results (that is, the regularity of minimizers up to a subset of the set of definition and the study of the properties of the singular set; see for example Section 4.2 in [1] and Section 1 in [2]) are contained in [3–10]. Only few everywhere regularity results are available: [11] where the everywhere continuity is proved in the two-dimensional case, [12] where Hölder continuity for extremals is dealt with in dimension two, [13] where local boundedness is proved in the three-dimensional case. Global pointwise bounds are in [14–19]. Interesting results are contained in [20–25]; see also [26, 27]. Let us come back to [13]; in such a paper, the authors make an important step toward regularity: they prove boundedness of minimizers in the three-dimensional case; unfortunately, they make restrictions that rule out the most important polyconvex integral. In the present paper, we find a different set of assumptions, which allows us to deal with such a polyconvex integral. In the next section, we write assumptions and results; in Sect. 3 we collect some preliminaries and, in Sect. 4, we give the proof of the main theorem.

2 Assumptions and Results

In this paper we study the regularity of vectorial local minimizers of integral functionals

$$I(v, \Omega) = \int_{\Omega} f(x, Dv(x)) dx, \quad (1)$$

where $\Omega \subset \mathbb{R}^3$ is an open, bounded set, $v : \Omega \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $v = (v^1, v^2, v^3)$ and Dv is the Jacobian matrix of its partial derivatives

$$Dv = (v_{x_i}^{\alpha})_{i=1,2,3}^{\alpha=1,2,3} = \begin{pmatrix} Dv^1 \\ Dv^2 \\ Dv^3 \end{pmatrix} = \begin{pmatrix} v_{x_1}^1 & v_{x_2}^1 & v_{x_3}^1 \\ v_{x_1}^2 & v_{x_2}^2 & v_{x_3}^2 \\ v_{x_1}^3 & v_{x_2}^3 & v_{x_3}^3 \end{pmatrix},$$

moreover, $f : \Omega \times \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty[$ is a Carathéodory function such that for fixed x

$$\xi \rightarrow f(x, \xi) \text{ is polyconvex}$$

that is

$$f(x, \xi) = g(x, \xi, \text{adj}_2 \xi, \det \xi) \quad \text{with} \quad (\xi, \lambda, t) \rightarrow g(x, \xi, \lambda, t) \quad \text{convex}, \quad (2)$$

see [28, 29]. When dealing with models in nonlinear elasticity, f is the stored-energy function; moreover, ξ , $\text{adj}_2 \xi$, $\det \xi$ govern the deformation of line, surface and volume elements respectively. Our model is

$$f(x, Dv) = |Dv|^p + |\text{adj}_2 Dv|^q + |\det Dv|^r, \quad (3)$$

where $\det Dv$ is the determinant of the matrix Dv , and $\text{adj}_2 Dv$ denotes the adjugate matrix of order 2, whose components are

$$(\text{adj}_2 Dv)_{ij} = (-1)^{i+j} \det \begin{pmatrix} v_{x_k}^\alpha & v_{x_\ell}^\alpha \\ v_{x_k}^\beta & v_{x_\ell}^\beta \end{pmatrix}, \quad i, j \in \{1, 2, 3\},$$

with $\alpha, \beta \in \{1, 2, 3\} \setminus \{i\}$, $\alpha < \beta$, and $k, \ell \in \{1, 2, 3\} \setminus \{j\}$, $k < \ell$. Moreover, $(\text{adj}_2 Dv)^\alpha$ denotes the α -row of $\text{adj}_2 Dv$, that is

$$(\text{adj}_2 Dv)^\alpha = ((\text{adj}_2 Dv)_{\alpha 1}, (\text{adj}_2 Dv)_{\alpha 2}, (\text{adj}_2 Dv)_{\alpha 3}).$$

In paper [13], the authors consider densities f for which the following splitting holds true

$$f(x, Dv) = \sum_{\alpha=1}^3 F^\alpha(x, Dv^\alpha) + \sum_{\beta=1}^3 G^\beta(x, (\text{adj}_2 Dv)^\beta) + H(x, \det Dv) \quad (4)$$

for suitable nonnegative functions F^α, G^β, H . Note that model (3), with $p \neq 2$, cannot be written as (4); see Lemma A.1 in “Appendix A”. In this paper, we succeed in dealing with model (3) and we prove the following

Theorem 2.1 *Let Ω be a bounded and open subset of \mathbb{R}^3 . Assume that $1 \leq r < q < p \leq 3$ with $2 < p$ and*

$$\begin{aligned} \frac{p}{p^*} &< \min \left\{ 1 - \frac{qp^*}{p(p^* - q)}, 1 - \frac{rp^*}{q(p^* - r)} \right\}, \quad \text{if } 1 < q \leq 2, \\ \frac{p}{p^*} &< \min \left\{ 1 - \frac{2p^*}{p(p^* - 2)} - \frac{(q-2)p^*}{q(p^* - 2)}, 1 - \frac{rp^*}{q(p^* - r)} \right\}, \quad \text{if } 2 < q; \end{aligned} \quad (5)$$

then all the local minimizers $u \in W_{loc}^{1,p}(\Omega; \mathbb{R}^3)$ of

$$\int_{\Omega} (|Du|^p + |\text{adj}_2 Du|^q + |\det Du|^r) \quad (6)$$

are locally bounded in Ω .

We recall that p^* is the Sobolev exponent: $p^* = \frac{np}{n-p} = \frac{3p}{3-p}$ when $p < n = 3$; moreover, p^* is any number greater than p when $p = n = 3$, so it can be chosen large enough that (5) is satisfied by assuming only $1 \leq r < q < p$. We notice that we have restricted ourselves to the case $p \leq 3$ because, when $p > 3$, every function in $W_{loc}^{1,p}(\Omega)$ is trivially in $L_{loc}^\infty(\Omega)$ by the Sobolev theorem. Note that we have existence of minimizers for (6) when $2 \leq p$, $\frac{p}{p-1} \leq q$ and $1 < r$, provided a boundary datum $\bar{u} \in W^{1,p}(\Omega; \mathbb{R}^3)$, with finite energy, has been fixed; see Remark 8.32 (iii) in [29] and Theorem 3.1 in [13]. Condition (5) is satisfied, for example, when $p = \frac{14}{5}$, $q = 2$, $r = \frac{3}{2}$ and this gives us the following.

Corollary 2.1 *Let Ω be a bounded and open subset of \mathbb{R}^3 and let $u \in W_{loc}^{1, \frac{14}{5}}(\Omega; \mathbb{R}^3)$ be a local minimizer of*

$$\int_{\Omega} (|Du|^{\frac{14}{5}} + |\text{adj}_2 Du|^2 + |\det Du|^{\frac{3}{2}}); \quad (7)$$

then u is locally bounded in Ω .

In the framework of Corollary 2.1, we have $\frac{p}{p-1} = \frac{14}{9} < 2 = q$, so the existence of minimizers is guaranteed as in the previous lines. Theorem 2.1 is a particular case of a more general result. Let us note that model (3) suggests we assume the following structure

$$f(x, \xi) = F(x, |\xi|^2) + G(x, |\text{adj}_2 \xi|^2) + H(x, \det \xi), \quad (8)$$

where F , G and H are Carathéodory nonnegative functions. We assume p -growth with respect to ξ , q -growth with respect to $\text{adj}_2 \xi$ and r -growth with respect to $\det \xi$

$$k_1 t^{p/2} - k_2 \leq F(x, t) \leq k_3 t^{p/2} + a(x) \quad (9)$$

$$k_1 t^{q/2} - k_2 \leq G(x, t) \leq k_3 t^{q/2} + b(x) \quad (10)$$

$$0 \leq H(x, s) \leq k_3 |s|^r + c(x), \quad (11)$$

where k_1, k_2, k_3 are constants such that $k_1, k_3 \in]0, +\infty[$ and $k_2 \in [0, +\infty[$ and $a, b, c : \Omega \rightarrow [0, +\infty[$ are functions in $L^\sigma(\Omega)$, $\sigma > 1$; as far as exponents p, q, r are concerned, we assume that $2 < p \leq 3$ and $1 \leq r < q < p$. Now we need to control the behavior of F with respect to the sum from below

$$F(x, t_1) + F(x, t_2) - k_2 \leq F(x, t_1 + t_2). \quad (12)$$

A weaker condition is needed for G :

$$G(x, t_1) - k_2 \leq G(x, t_1 + t_2). \quad (13)$$

We also need to control the behavior of F with respect to the sum from above:

$$F(x, t_1 + t_2) \leq F(x, t_1) + F(x, t_2) + k_3 t_1 t_2^{\frac{p}{2}-1} + a(x). \quad (14)$$

Note that in (14) there is an extra term with the product between t_1 and t_2 . When $q > 2$ we assume

$$G(x, t_1 + t_2) \leq G(x, t_1) + G(x, t_2) + k_3 t_1 t_2^{\frac{q}{2}-1} + b(x). \quad (15)$$

When $q \leq 2$ we do not need the product between t_1 and t_2 any longer; we require subadditivity

$$G(x, t_1 + t_2) \leq G(x, t_1) + G(x, t_2) + b(x). \quad (16)$$

Functions F verifying the previous assumptions are $F(x, t) = \gamma(x)t^{p/2}$ and $F(x, t) = \gamma(x)(1 + t^2)^{p/4}$, provided $\gamma(x)$ is positive and away from both 0 and $+\infty$; similar examples for G and H : see Remarks 3.2, ..., 3.7. Our main result is the following

Theorem 2.2 *Let Ω be a bounded and open subset of \mathbb{R}^3 and let f be as in (8); assume that conditions (9)–(16) hold with $1 \leq r < q < p \leq 3$ such that $2 < p$ and*

$$\begin{aligned} \frac{p}{p^*} &< \min \left\{ 1 - \frac{qp^*}{p(p^* - q)}, 1 - \frac{rp^*}{q(p^* - r)}, 1 - \frac{1}{\sigma} \right\}, \quad \text{if } 1 < q \leq 2, \\ \frac{p}{p^*} &< \min \left\{ 1 - \frac{2p^*}{p(p^* - 2)} - \frac{(q - 2)p^*}{q(p^* - 2)}, 1 - \frac{rp^*}{q(p^* - r)}, 1 - \frac{1}{\sigma} \right\}, \quad \text{if } 2 < q. \end{aligned} \quad (17)$$

Then, all the local minimizers $u \in W_{loc}^{1,p}(\Omega; \mathbb{R}^3)$ of I are locally bounded in Ω .

Note that $\frac{1}{\sigma} = 0$, if $\sigma = \infty$. In our Theorem 2.2, we assume (8); in [13] (4) was in force: in vectorial problems, some structure conditions are due to minimizers which can be unbounded: see De Giorgi's counterexample [30]; see also [31], Section 3 in [1] and [32]. As far as exponents p, q, r are concerned, (17) is the same as (2.5) in [13] when $1 < q \leq 2$; if $2 < q$ then (17) seems to require a bit more than (2.5) in [13]: see comparison (72).

The integrals we consider show a \tilde{p} growth from below and a \tilde{q} growth from above, so we are in the class of functionals with \tilde{p}, \tilde{q} -growth. It is now well known, as in our result, that a restriction between \tilde{p} and \tilde{q} must be imposed due to counterexamples in [33–37]; see also [38, 39]; we refer to [1] for a detailed survey on the subject.

3 Preliminaries

In this section, we recall some standard definitions and collect several lemmas useful in our proofs.

First of all, we recall the following

Definition 3.1 A function $u \in W_{loc}^{1,1}(\Omega; \mathbb{R}^3)$ is a local minimizer of (1) if $f(Du) \in L_{loc}^1(\Omega)$ and

$$I(u, \text{supp } \varphi) \leq I(u + \varphi, \text{supp } \varphi), \quad (18)$$

for all $\varphi \in W^{1,1}(\Omega, \mathbb{R}^3)$ with $\text{supp } \varphi \subset \subset \Omega$.

All the norms we use on \mathbb{R}^3 and $\mathbb{R}^{3 \times 3}$ will be the standard Euclidean ones and denoted by $|\cdot|$ in all cases. In particular, for matrices $\xi, \eta \in \mathbb{R}^{3 \times 3}$ we write $\langle \xi, \eta \rangle := \text{trace}(\xi^T \eta)$ for the usual inner product of ξ and η , and $|\xi| := \langle \xi, \xi \rangle^{\frac{1}{2}}$ for the corresponding Euclidean norm.

Lemma 3.1 *For $a, b \geq 0$ we have that*

$$a^m + b^m \leq (a + b)^m, \quad \text{if } m \geq 1, \quad (19)$$

$$(a + b)^m \leq a^m + b^m + mab^{m-1}, \quad \text{if } 1 \leq m \leq 2. \quad (20)$$

Proof When $m = 1$, (19) and (20) are easy. We are left with the case $1 < m$. It is obvious that (19) and (20) hold true both for $b = 0$. We now assume $b > 0$ and we let $t = a/b$. It suffices to show that for $t \geq 0$,

$$t^m + 1 \leq (t + 1)^m, \quad \text{if } m > 1, \quad (21)$$

$$(t + 1)^m \leq t^m + 1 + mt, \quad \text{if } 1 < m \leq 2. \quad (22)$$

In order to prove (21), we let $h(t) = (t + 1)^m - t^m - 1$. Since

$$h(0) = 0 \quad (23)$$

and, by $m > 1$,

$$h'(t) = m[(t + 1)^{m-1} - t^{m-1}] \geq 0, \quad (24)$$

then $h(t) \geq 0$ and (21) follows.

Regarding (22), we let $g(t) = (t + 1)^m - t^m - mt - 1$. Since

$$g(0) = 0, \quad (25)$$

$$g'(t) = m[(t + 1)^{m-1} - t^{m-1} - 1] \leq m[t^{m-1} + 1 - t^{m-1} - 1] = 0, \quad (26)$$

where we used $1 < m \leq 2$ and Remark 3.1, then (22) follows. \square

Remark 3.1 We recall the well-known inequality: for $a, b \geq 0$ we have

$$(a + b)^m \leq a^m + b^m, \quad \text{if } 0 < m \leq 1. \quad (27)$$

Lemma 3.2 Fix $m \in [-\frac{1}{2}, +\infty[$ and consider $V : \mathbb{R} \rightarrow \mathbb{R}$ as follows

$$V(s) = \left(1 + s^2\right)^m s; \quad (28)$$

then, $V : \mathbb{R} \rightarrow \mathbb{R}$ is strictly increasing.

Proof We compute the first derivative

$$V'(s) = \left(1 + s^2\right)^{m-1} [(2m + 1)s^2 + 1]; \quad (29)$$

since $m \geq -\frac{1}{2}$, we have $V'(s) > 0$ for every $s \in \mathbb{R}$. This ends the proof. \square

Lemma 3.3 Fix $p \in [2, +\infty[$; consider $w : \mathbb{R}^2 \rightarrow \mathbb{R}$ as follows

$$w(a, b) = [1 + (a + b)^2]^{p/4} - [1 + a^2]^{p/4} - [1 + b^2]^{p/4}. \quad (30)$$

Then,

$$a \geq 0, \quad b \geq 0 \implies w(0, 0) \leq w(a, b). \quad (31)$$

Proof We compute the first partial derivatives:

$$\begin{aligned}\frac{\partial w}{\partial a}(a, b) &= \frac{p}{4}[1 + (a + b)^2]^{(p/4)-1}2(a + b) - \frac{p}{4}[1 + a^2]^{(p/4)-1}2a \\ &= \frac{p}{2}\{V(a + b) - V(a)\}\end{aligned}$$

and

$$\begin{aligned}\frac{\partial w}{\partial b}(a, b) &= \frac{p}{4}[1 + (a + b)^2]^{(p/4)-1}2(a + b) - \frac{p}{4}[1 + b^2]^{(p/4)-1}2b \\ &= \frac{p}{2}\{V(a + b) - V(b)\}\end{aligned}$$

where V is given by (28) with $m = (p/4) - 1$. Note that $m \geq -1/2$ since $p \geq 2$. Then, V is increasing so that, when $a \geq 0$ and $b \geq 0$, we have $V(a + b) - V(a) \geq 0$ and $V(a + b) - V(b) \geq 0$. This shows that

$$a \geq 0, \quad b \geq 0 \implies \frac{\partial w}{\partial a}(a, b) \geq 0, \quad \frac{\partial w}{\partial b}(a, b) \geq 0. \quad (32)$$

Then, $a \rightarrow w(a, b)$ increases and $b \rightarrow w(a, b)$ increases too, if we restrict ourselves to $a \geq 0$ and $b \geq 0$; thus,

$$w(0, 0) \leq w(0, b) \leq w(a, b), \quad (33)$$

provided $b \geq 0$ and $a \geq 0$. This ends the proof. \square

Corollary 3.1 Fix $p \in [2, +\infty[$; then,

$$a \geq 0, \quad b \geq 0 \implies [1 + a^2]^{p/4} + [1 + b^2]^{p/4} - 1 \leq [1 + (a + b)^2]^{p/4}. \quad (34)$$

Proof We write (31) explicitly and we get (34). \square

Lemma 3.4 Fix $p \in]2, 3]$. If $a \geq 0$ and $b \geq 0$, then

$$[1 + (a + b)^2]^{p/4} \leq [1 + a^2]^{p/4} + [1 + b^2]^{p/4} + \frac{p}{2}ab^{(p/2)-1} + 1. \quad (35)$$

Proof Since $p \in]2, 3]$ we have $\frac{p}{4} \in [\frac{1}{2}, \frac{3}{4}]$ and we can use (27) with $m = \frac{p}{4}$:

$$[1 + (a + b)^2]^{p/4} \leq [1]^{p/4} + [(a + b)^2]^{p/4} = 1 + (a + b)^{p/2}; \quad (36)$$

now $\frac{p}{2} \in]1, \frac{3}{2}]$ and we can use (20) with $m = \frac{p}{2}$:

$$\begin{aligned} 1 + (a + b)^{p/2} &\leq 1 + a^{p/2} + b^{p/2} + \frac{p}{2} ab^{(p/2)-1} \\ &= 1 + (a^2)^{p/4} + (b^2)^{p/4} + \frac{p}{2} ab^{(p/2)-1} \\ &\leq 1 + [1 + a^2]^{p/4} + [1 + b^2]^{p/4} + \frac{p}{2} ab^{(p/2)-1}. \end{aligned} \quad (37)$$

This ends the proof. \square

Lemma 3.5 Fix $q \in]1, 2]$. Then,

$$a \geq 0, \quad b \geq 0 \implies [1 + (a + b)^2]^{q/4} \leq [1 + a^2]^{q/4} + [1 + b^2]^{q/4} + 1. \quad (38)$$

Proof Since $q \in]1, 2]$ we have $\frac{q}{4} \in [\frac{1}{4}, \frac{1}{2}]$ and we can use (27) with $m = \frac{q}{4}$:

$$[1 + (a + b)^2]^{q/4} \leq 1^{q/4} + [(a + b)^2]^{q/4} = 1 + (a + b)^{q/2}; \quad (39)$$

now $\frac{q}{2} \in [\frac{1}{2}, 1]$ and we can use (27) with $m = \frac{q}{2}$:

$$\begin{aligned} 1 + (a + b)^{q/2} &\leq 1 + a^{q/2} + b^{q/2} \\ &= 1 + (a^2)^{q/4} + (b^2)^{q/4} \\ &\leq 1 + [1 + a^2]^{q/4} + [1 + b^2]^{q/4}. \end{aligned} \quad (40)$$

This ends the proof. \square

Now we are able to give examples of functions F, G, H verifying conditions required in Theorem 2.2.

Remark 3.2 Fix $p \in]2, 3]$ and define

$$F(x, t) = \gamma(x)t^{p/2} \quad (41)$$

for $t \in [0, +\infty[$, where $\gamma_1 \leq \gamma(x) \leq \gamma_2$ with $\gamma_1, \gamma_2 \in]0, +\infty[$. Then (9), (12), (14) hold true with $k_1 = \gamma_1, k_2 = 0, k_3 = \frac{p}{2}\gamma_2, a(x) = 0$. Indeed, we use (19) and (20) with $m = p/2$ in Lemma 3.1 and we are done.

Remark 3.3 Fix $q \in]1, 3[$ and define

$$G(x, t) = \gamma(x)t^{q/2} \quad (42)$$

for $t \in [0, +\infty[$, where $\gamma_1 \leq \gamma(x) \leq \gamma_2$ with $\gamma_1, \gamma_2 \in]0, +\infty[$. Then, when $q > 2$, (10), (13), (15) hold true with $k_1 = \gamma_1, k_2 = 0, k_3 = \frac{q}{2}\gamma_2, b(x) = 0$. Indeed, we use (20) with $m = q/2$ in Lemma 3.1 and we are done. Moreover, when $q \leq 2$, (10), (13), (16) hold true with $k_1 = \gamma_1, k_2 = 0, k_3 = \gamma_2, b(x) = 0$. Indeed, when $q \leq 2$, we use the well-known inequality (27) with $m = q/2$ and we are done.

Remark 3.4 Fix $r \in [1, 3[$ and define

$$H(x, s) = \gamma(x)|s|^r \quad (43)$$

for $s \in \mathbb{R}$, where $\gamma_1 \leq \gamma(x) \leq \gamma_2$ with $\gamma_1, \gamma_2 \in]0, +\infty[$. Then, (11) holds true with $k_3 = \gamma_2, c(x) = 0$.

Remark 3.5 Fix $p \in]2, 3]$ and define

$$F(x, t) = \gamma(x)[1 + t^2]^{p/4} \quad (44)$$

for $t \in [0, +\infty[$, where $\gamma_1 \leq \gamma(x) \leq \gamma_2$ with $\gamma_1, \gamma_2 \in]0, +\infty[$. Then (9), (12), (14) hold true with $k_1 = \gamma_1, k_2 = \gamma_2, k_3 = \frac{p}{2}\gamma_2, a(x) = \gamma_2$. Indeed, we use (27) with $m = p/4$, (34) and (35).

Remark 3.6 Fix $q \in]1, 3[$ and define

$$G(x, t) = \gamma(x)[1 + t^2]^{q/4} \quad (45)$$

for $t \in [0, +\infty[$, where $\gamma_1 \leq \gamma(x) \leq \gamma_2$ with $\gamma_1, \gamma_2 \in]0, +\infty[$. Then, when $q > 2$, (10), (13), (15) hold true with $k_1 = \gamma_1, k_2 = 0, k_3 = \frac{q}{2}\gamma_2, b(x) = \gamma_2$. Indeed, we use (27) with $m = q/4$, (35) and we are done. Moreover, when $q \leq 2$, (10), (13), (16) hold true with $k_1 = \gamma_1, k_2 = 0, k_3 = \gamma_2, b(x) = \gamma_2$. Indeed, when $q \leq 2$, we use (27) with $m = q/4$, (38) and we are done.

Remark 3.7 Fix $r \in [1, 3[$ and define

$$H(x, s) = \gamma(x)[1 + |s|^2]^{r/2} \quad (46)$$

for $s \in \mathbb{R}$, where $\gamma_1 \leq \gamma(x) \leq \gamma_2$ with $\gamma_1, \gamma_2 \in]0, +\infty[$. Then, (11) holds true with $k_3 = 2^{r/2}\gamma_2, c(x) = 2^{r/2}\gamma_2$.

The following lemma can be found in [13] as Lemma 4.1.

Lemma 3.6 Consider the matrices $A, B \in \mathbb{R}^{3 \times 3}$:

$$A = \begin{pmatrix} A^1 \\ B^2 \\ B^3 \end{pmatrix}, \quad B = \begin{pmatrix} B^1 \\ B^2 \\ B^3 \end{pmatrix}.$$

Then, the following estimates hold:

- (a) $|A| \leq |A^1| + |B^2| + |B^3|$,
- (b) $|\det A| \leq |A^1| |(\text{adj}_2 B)^1|$,
- (c) $|(\text{adj}_2 A)_{2j}| \leq |A^1| |B^3|$ and $|(\text{adj}_2 A)_{3j}| \leq |A^1| |B^2|$, for all $j \in \{1, 2, 3\}$.

In order to get our main result, we have to prove a suitable Caccioppoli-type inequality for any component u^α of the local minimizer u of functional I (1) on every superlevel set $\{u^\alpha > k\}$. To this goal, we will use the following lemma (see [13] for a proof).

Lemma 3.7 *Let Ω be an open subset of \mathbb{R}^3 . Consider a Carathéodory function $f : \Omega \times \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty]$. Assume that there exist $c_1, c_3 > 0$ and $c_2 \geq 0$ such that, for every $\xi \in \mathbb{R}^{3 \times 3}$,*

$$c_1(|\xi|^p + |\text{adj}_2 \xi|^q) - c_2 \leq f(x, \xi) \leq c_3(|\xi|^p + |(\text{adj}_2 \xi)|^q + |\det \xi|^r + 1 + \omega(x)),$$

with $1 \leq p, 1 \leq q, 1 \leq r, \omega(x) \geq 0$.

Let $u \in W_{loc}^{1,p}(\Omega; \mathbb{R}^3)$ be such that $x \rightarrow f(x, Du(x)) \in L_{loc}^1(\Omega)$. Fix $\eta \in C_0^1(\Omega)$, $\eta \geq 0$ and $k \in \mathbb{R}$, and denote, for almost every $x \in \{u^1 > k\} \cap \{\eta > 0\}$,

$$A = \begin{pmatrix} \mu \eta^{-1}(k - u^1) D\eta \\ Du^2 \\ Du^3 \end{pmatrix}.$$

If

$$q < \frac{p^* p}{p^* + p} \quad \text{and} \quad r < \frac{p^* q}{p^* + q}$$

and $\omega \in L_{loc}^1(\Omega)$, then

$$\eta^\mu f(x, A) \in L^1(\{u^1 > k\} \cap \{\eta > 0\}), \quad \forall \mu \geq p^*.$$

4 Proof of Theorem 2.2

We want to stress that the proof of our result follows the idea used in [13]: we provide the local boundedness of the minimizers by proving that each component is locally bounded. In the following lemma, we refer to the first component u^1 : the core of the proof lies in the following Caccioppoli-type inequality, obtained on every superlevel set $\{u^1 > k\}$. We keep in mind that $p^* = \frac{np}{n-p}$ if $p < n = 3$ and p^* is any number $> p$ when $p = n = 3$.

Proposition 4.1 (Caccioppoli-type estimate) *Let f be as in (8) satisfying (9)–(16) with $1 \leq r < q < p \leq 3$ such that*

$$2 < p, \quad q < \frac{pp^*}{p + p^*}, \quad r < \frac{p^* q}{p^* + q}. \quad (47)$$

Let $u \in W_{loc}^{1,p}(\Omega; \mathbb{R}^3)$ be a local minimizer of I . Let $B_R(x_0) \subset \subset \Omega$ with $|B_R(x_0)| < 1$; fixed $k \in \mathbb{R}$, denote

$$A_{k,\tau}^1 := \{x \in B_\tau(x_0) : u^1(x) > k\} \quad 0 < \tau \leq R.$$

Then, there exists $C = C(k_1, k_2, k_3, p, q, r, p^) > 0$ such that, for every $0 < s < t \leq R$:*

$$\begin{aligned}
 \int_{A_{k,s}^1} |Du^1|^p dx &\leq C \int_{A_{k,t}^1} \left(\frac{u^1 - k}{t - s} \right)^{p^*} dx + C \left\{ 1 + \|a + b + c\|_{L^\sigma(B_R)} \right. \\
 &+ \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{p^*(p-2)}{p(p^*-2)}} + \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{qp^*}{p(p^*-q)}} \\
 &+ \left(\int_{B_R} |(\text{adj}_2 Du)^1|^q dx \right)^{\frac{rp^*}{q(p^*-r)}} + 1_{(2,+\infty)}(q) \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{2p^*}{p(p^*-2)}} \\
 &\times \left. \left(\int_{B_R} |(\text{adj}_2 Du)^1|^q dx \right)^{\frac{(q-2)p^*}{q(p^*-2)}} \right\} |A_{k,t}^1|^\theta,
 \end{aligned} \tag{48}$$

where

$$\begin{aligned}
 \theta &:= \min \left\{ 1 - \frac{qp^*}{p(p^*-q)}, 1 - \frac{rp^*}{q(p^*-r)}, 1 - \frac{1}{\sigma} \right\}, \quad \text{if } 1 < q \leq 2, \\
 \theta &:= \min \left\{ 1 - \frac{2p^*}{p(p^*-2)} - \frac{(q-2)p^*}{q(p^*-2)}, 1 - \frac{rp^*}{q(p^*-r)}, 1 - \frac{1}{\sigma} \right\}, \quad \text{if } 2 < q,
 \end{aligned}$$

with $\frac{1}{\sigma} = 0$ if $\sigma = \infty$.

Proof The condition $|B_R(x_0)| = \frac{4\pi R^3}{3} < 1$ ensures $R < 1$. Let s, t be such that $0 < s < t \leq R$. Consider a cutoff function $\eta \in C_0^\infty(B_t(x_0))$ satisfying the following assumptions:

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ in } B_s(x_0), \quad |D\eta| \leq \frac{2}{t-s}.$$

Fixing $k \in \mathbb{R}$, define $w \in W_{loc}^{1,p}(\Omega; \mathbb{R}^3)$,

$$w^1 := \max\{u^1 - k, 0\}, \quad w^2 = 0, \quad w^3 = 0,$$

and, for $\mu = p^*$,

$$\varphi := -\eta^\mu w.$$

For almost every $x \in \Omega \setminus (\{\eta > 0\} \cap \{u^1 > k\})$ we have $\varphi = 0$, thus

$$f(x, Du + D\varphi) = f(x, Du) \tag{49}$$

almost everywhere in $\Omega \setminus (\{\eta > 0\} \cap \{u^1 > k\})$.

For almost every $x \in \{\eta > 0\} \cap \{u^1 > k\}$ denote

$$A = \begin{pmatrix} \mu\eta^{-1}(k - u^1)D\eta \\ Du^2 \\ Du^3 \end{pmatrix}. \tag{50}$$

We notice that

$$Du + D\varphi = \begin{pmatrix} (1 - \eta^\mu)Du^1 + \mu\eta^{\mu-1}(k - u^1)D\eta \\ Du^2 \\ Du^3 \end{pmatrix} = (1 - \eta^\mu)Du + \eta^\mu A.$$

Moreover, since for almost every $x \in \{\eta > 0\} \cap \{u^1 > k\}$,

$$\det(Du + D\varphi) = (1 - \eta^\mu) \det Du + \eta^\mu \det A$$

and

$$\text{adj}_2(Du + D\varphi) = (1 - \eta^\mu)\text{adj}_2 Du + \eta^\mu \text{adj}_2 A,$$

then, since f is polyconvex, we get that

$$f(x, Du + D\varphi) \leq (1 - \eta^\mu)f(x, Du) + \eta^\mu f(x, A) \quad (51)$$

almost everywhere in $\{\eta > 0\} \cap \{u^1 > k\}$.

By the minimality of u , $f(x, Du) \in L^1_{loc}(\Omega)$; note that in our case we can use Lemma 3.7, deducing that

$$\eta^\mu f(x, A) \in L^1(\{\eta > 0\} \cap \{u^1 > k\}).$$

Therefore, (49) and (51) imply $f(x, Du + D\varphi) \in L^1_{loc}(\Omega)$.

By the local minimality of u , (49) and (51), recalling that $A^1_{k,t}$ is the set $\{x \in B_t(x_0) : u^1(x) > k\}$, we have

$$\begin{aligned} \int_{A^1_{k,t} \cap \{\eta > 0\}} f(x, Du) dx &\leq \int_{A^1_{k,t} \cap \{\eta > 0\}} f(x, Du + D\varphi) dx \\ &\leq \int_{A^1_{k,t} \cap \{\eta > 0\}} \{(1 - \eta^\mu)f(x, Du) + \eta^\mu f(x, A)\} dx. \end{aligned}$$

The inequality above implies

$$\int_{A^1_{k,t} \cap \{\eta > 0\}} \eta^\mu f(x, Du) dx \leq \int_{A^1_{k,t} \cap \{\eta > 0\}} \eta^\mu f(x, A) dx.$$

Taking into account the expression of f (see (8)), we obtain from the above inequality that

$$\begin{aligned} &\int_{A^1_{k,t} \cap \{\eta > 0\}} \eta^\mu \left[F(x, |Du|^2) + G(x, |\text{adj}_2 Du|^2) + H(x, \det Du) \right] dx \\ &\leq \int_{A^1_{k,t} \cap \{\eta > 0\}} \eta^\mu \left[F(x, |A|^2) + G(x, |\text{adj}_2 A|^2) + H(x, \det A) \right] dx. \quad (52) \end{aligned}$$

Denote $\tilde{u} = (u^2, u^3)$ and

$$D\tilde{u} = \begin{pmatrix} Du^2 \\ Du^3 \end{pmatrix}.$$

We have

$$|Du|^2 = |Du^1|^2 + |D\tilde{u}|^2;$$

we use (12) with $t_1 = |Du^1|^2$ and $t_2 = |D\tilde{u}|^2$, so that

$$\begin{aligned} & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[F(x, |Du^1|^2) + F(x, |D\tilde{u}|^2) - k_2 \right] dx \\ & \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu F(x, |Du|^2) dx. \end{aligned} \quad (53)$$

Note that

$$|A|^2 = |A^1|^2 + |D\tilde{u}|^2;$$

by using (14) with $t_1 = |A^1|^2$ and $t_2 = |D\tilde{u}|^2$, we obtain

$$\begin{aligned} & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu F(x, |A|^2) dx \\ & \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[F(x, |A^1|^2) + F(x, |D\tilde{u}|^2) \right. \\ & \quad \left. + k_3 |A^1|^2 (|D\tilde{u}|^2)^{\frac{p}{2}-1} + a(x) \right] dx. \end{aligned} \quad (54)$$

Furthermore, setting

$$|\tilde{A}|^2 = |(\text{adj}_2 Du)^2|^2 + |(\text{adj}_2 Du)^3|^2, \quad |\tilde{\tilde{A}}|^2 = |(\text{adj}_2 A)^2|^2 + |(\text{adj}_2 A)^3|^2 \quad (55)$$

and noticing

$$(\text{adj}_2 A)^1 = (\text{adj}_2 Du)^1,$$

we can write

$$\begin{aligned} |\text{adj}_2 Du|^2 &= |(\text{adj}_2 Du)^1|^2 + |(\text{adj}_2 Du)^2|^2 + |(\text{adj}_2 Du)^3|^2 = |(\text{adj}_2 Du)^1|^2 + |\tilde{A}|^2, \\ |\text{adj}_2 A|^2 &= |(\text{adj}_2 A)^1|^2 + |(\text{adj}_2 A)^2|^2 + |(\text{adj}_2 A)^3|^2 = |(\text{adj}_2 Du)^1|^2 + |\tilde{\tilde{A}}|^2. \end{aligned}$$

Applying (13) with $t_1 = |(\text{adj}_2 Du)^1|^2$ and $t_2 = |\tilde{A}|^2$, we get

$$\begin{aligned} & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left(G(x, |(\text{adj}_2 Du)^1|^2) - k_2 \right) dx \\ & \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu G(x, |\text{adj}_2 Du|^2) dx. \end{aligned} \quad (56)$$

Assumption (15) when $q > 2$ or (16) when $q \leq 2$, with $t_1 = |\tilde{A}|^2$ and $t_2 = |(\text{adj}_2 Du)^1|^2$, yields

$$\begin{aligned} & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu G(x, |\text{adj}_2 A|^2) dx \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu [G(x, |\tilde{A}|^2) \\ & + G(x, |(\text{adj}_2 Du)^1|^2) + b(x) + 1_{(2,+\infty)}(q) k_3 |\tilde{A}|^2 \left(|(\text{adj}_2 Du)^1|^2 \right)^{\frac{q}{2}-1}] dx. \end{aligned} \quad (57)$$

By virtue of (53), (54), (56) and (57), from (52), we get

$$\begin{aligned} & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu [F(x, |Du^1|^2) + F(x, |D\tilde{u}|^2) - 2k_2 \\ & + G(x, |(\text{adj}_2 Du)^1|^2) + H(x, \det Du)] dx \\ & \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[F(x, |A^1|^2) + F(x, |D\tilde{u}|^2) + k_3 |A^1|^2 (|D\tilde{u}|^2)^{\frac{p}{2}-1} \right. \\ & + a(x) + G(x, |\tilde{A}|^2) + G(x, |(\text{adj}_2 Du)^1|^2) + b(x) \\ & \left. + 1_{(2,+\infty)}(q) k_3 |\tilde{A}|^2 \left(|(\text{adj}_2 Du)^1|^2 \right)^{\frac{q}{2}-1} + H(x, \det A) \right] dx \end{aligned}$$

and then

$$\begin{aligned} & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[F(x, |Du^1|^2) - 2k_2 + H(x, \det Du) \right] dx \\ & \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[F(x, |A^1|^2) + k_3 |A^1|^2 (|D\tilde{u}|^2)^{\frac{p}{2}-1} + a(x) + G(x, |\tilde{A}|^2) \right. \\ & \left. + b(x) + 1_{(2,+\infty)}(q) k_3 |\tilde{A}|^2 \left(|(\text{adj}_2 Du)^1|^2 \right)^{\frac{q}{2}-1} + H(x, \det A) \right] dx. \end{aligned} \quad (58)$$

In order to estimate the first two terms on the right-hand side of (58), we recall that $\mu = p^* > p$ and

$$A^1 = \mu \eta^{-1} (k - u^1) D\eta.$$

By using the right-hand side of (9) and the fact $z^p \leq 1 + z^{p^*}$ if $z \geq 0$, we obtain

$$\begin{aligned}
 \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu F(x, |A^1|^2) &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[k_3 |A^1|^p + a(x) \right] dx \\
 &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left[k_3 \left\{ 1 + |A^1|^{p^*} \right\} + a(x) \right] dx \\
 &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \left\{ \eta^\mu (k_3 + a(x)) + k_3 (2\mu)^{p^*} \eta^{\mu-p^*} \left(\frac{u^1 - k}{t-s} \right)^{p^*} \right\} dx \\
 &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \left\{ k_3 + a(x) + k_3 (2\mu)^{p^*} \left(\frac{u^1 - k}{t-s} \right)^{p^*} \right\} dx. \tag{59}
 \end{aligned}$$

We will write d' to denote the Hölder conjugate of $d > 1$: $d' = \frac{d}{d-1}$. Regarding the second term on the right-hand side in (58), notice that

$$(p-2) \left(\frac{p^*}{2} \right)' < p;$$

we use Young inequality with $\frac{p^*}{2}$, $\left(\frac{p^*}{2} \right)'$ and Hölder inequality with $\frac{p}{(p-2)\left(\frac{p^*}{2} \right)'}$, $\frac{p}{p-(p-2)\left(\frac{p^*}{2} \right)'}$:

$$\begin{aligned}
 &\int_{A_{k,t}^1 \cap \{\eta > 0\}} k_3 \eta^\mu |A^1|^2 |D\tilde{u}|^{p-2} dx \\
 &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} k_3 \eta^\mu |A^1|^{p^*} dx + \int_{A_{k,t}^1 \cap \{\eta > 0\}} k_3 \eta^\mu |D\tilde{u}|^{(p-2)\left(\frac{p^*}{2} \right)'} dx \\
 &\leq k_3 (2\mu)^{p^*} \int_{A_{k,t}^1 \cap \{\eta > 0\}} \left(\frac{u^1 - k}{t-s} \right)^{p^*} dx \\
 &\quad + k_3 \left(\int_{B_R} |D\tilde{u}|^p dx \right)^{\left(1 - \frac{2}{p} \right) \left(\frac{p^*}{2} \right)'} |A_{k,t}^1|^{1 - \left(1 - \frac{2}{p} \right) \left(\frac{p^*}{2} \right)'}. \tag{60}
 \end{aligned}$$

Now we estimate the fourth term in (58). By using (10), (55), Lemma 3.6-(c) and Young inequality with exponents $\frac{p^*}{q}$ and $\left(\frac{p^*}{q} \right)'$, we estimate

$$\begin{aligned}
 \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu G(x, |\tilde{A}|^2) &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu k_3 \left((|\tilde{A}|^2)^{\frac{q}{2}} + b(x) \right) dx \\
 &= \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left(k_3 \left(|(\text{adj}_2 A)^2|^2 + |(\text{adj}_2 A)^3|^2 \right)^{\frac{q}{2}} + b(x) \right) dx \\
 &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left(3^{\frac{q}{2}} k_3 \left[|A^1|^2 (|Du^2|^2 + |Du^3|^2) \right]^{\frac{q}{2}} + b(x) \right) dx
 \end{aligned}$$

$$\begin{aligned} &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu 3^{\frac{q}{2}} k_3 |A^1|^{p^*} dx + \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu 3^{\frac{q}{2}} k_3 (|D\tilde{u}|)^{q(\frac{p^*}{q})'} dx \\ &\quad + \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu b(x) dx. \end{aligned}$$

Note that $q(\frac{p^*}{q})' < p$ when $q < \frac{pp^*}{p+p^*}$ and $\frac{pp^*}{p+p^*} > 1$ if $p > \frac{2n}{n+1}$.

Moreover, Hölder inequality, with $\frac{p}{q} / \left(\frac{p^*}{q}\right)'$ and $\left(\frac{p}{q} / \left(\frac{p^*}{q}\right)'\right)'$, yields

$$\begin{aligned} \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |D\tilde{u}|^{q(\frac{p^*}{q})'} dx &\leq \left(\int_{A_{k,t}^1} (|D\tilde{u}|)^p dx \right)^{\frac{q}{p}(\frac{p^*}{q})'} |A_{k,t}^1|^{1-\frac{q}{p}(\frac{p^*}{q})'} \\ &\leq \left(\int_{B_R} (|D\tilde{u}|)^p dx \right)^{\frac{q}{p}(\frac{p^*}{q})'} |A_{k,t}^1|^{1-\frac{q}{p}(\frac{p^*}{q})'}; \end{aligned} \quad (61)$$

therefore, if we note that $(\frac{p^*}{q})' = \frac{p^*}{p^*-q}$, we have

$$\begin{aligned} \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu G(x, |\tilde{A}|^2) &\leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu 3^{\frac{q}{2}} k_3 |A^1|^{p^*} dx \\ &\quad + 3^{\frac{q}{2}} k_3 \left(\int_{B_R} |D\tilde{u}|^p dx \right)^{\frac{qp^*}{p(p^*-q)}} |A_{k,t}^1|^{1-\frac{qp^*}{p(p^*-q)}} + \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu b(x) dx. \end{aligned} \quad (62)$$

Eventually, if $q > 2$, we have to estimate the sixth term in (58) and we use Lemma 3.6-(c) and Young inequality with exponents $\frac{p^*}{2}$ and $\left(\frac{p^*}{2}\right)'$, so having

$$\begin{aligned} &\int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |\tilde{A}|^2 \left(|(\text{adj}_2 Du)^1|^2 \right)^{\frac{q}{2}-1} dx \\ &= \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu \left(|(\text{adj}_2 A)^2|^2 + |(\text{adj}_2 A)^3|^2 \right) |(\text{adj}_2 Du)^1|^{q-2} dx \\ &\leq 3 \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |A^1|^2 (|Du^3|^2 + |Du^2|^2) |(\text{adj}_2 Du)^1|^{q-2} dx \\ &\leq 3 \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |A^1|^{p^*} dx \\ &\quad + 3 \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |D\tilde{u}|^{2(\frac{p^*}{2})'} |(\text{adj}_2 Du)^1|^{(q-2)(\frac{p^*}{2})'} dx. \end{aligned} \quad (63)$$

Observe that $2 < q < \frac{pp^*}{p+p^*}$ implies $p > \frac{12}{5}$; so we have $2\left(\frac{p^*}{2}\right)' < p$ and we apply

Hölder inequality with exponents $\frac{p}{2(\frac{p^*}{2})'}$ and $\left(\frac{p}{2(\frac{p^*}{2})'}\right)'$,

$$\begin{aligned}
 & \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |D\tilde{u}|^{2\left(\frac{p^*}{2}\right)'} |(\text{adj}_2 Du)^1|^{(q-2)\left(\frac{p^*}{2}\right)'} dx \\
 & \leq \left(\int_{B_R} |D\tilde{u}|^p \right)^{\frac{2}{p}\left(\frac{p^*}{2}\right)'} \\
 & \quad \times \left(\int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |(\text{adj}_2 Du)^1|^{(q-2)\left(\frac{p^*}{2}\right)'} \left(\frac{p}{2\left(\frac{p^*}{2}\right)'} \right)' \right)^{1 - \frac{2}{p}\left(\frac{p^*}{2}\right)'} \quad (64)
 \end{aligned}$$

Furthermore, if $(q-2)\left(\frac{p^*}{2}\right)'\left(\frac{p}{2\left(\frac{p^*}{2}\right)'}\right)' < q$, we apply Hölder inequality again with exponents $\frac{q}{(q-2)\left(\frac{p^*}{2}\right)'\left(\frac{p}{2\left(\frac{p^*}{2}\right)'}\right)'}$ and its conjugate:

$$\begin{aligned}
 & \left(\int_{B_R} |D\tilde{u}|^p \right)^{\frac{2}{p}\left(\frac{p^*}{2}\right)'} \left(\int_{A_{k,t}^1} |(\text{adj}_2 Du)^1|^{(q-2)\left(\frac{p^*}{2}\right)'\left(\frac{p}{2\left(\frac{p^*}{2}\right)'}\right)'} \right)^{1 - \frac{2}{p}\left(\frac{p^*}{2}\right)'} \\
 & \leq \left(\int_{B_R} |D\tilde{u}|^p \right)^{\frac{2}{p}\left(\frac{p^*}{2}\right)'} \left[\left(\int_{A_{k,t}^1} |(\text{adj}_2 Du)^1|^q \right)^{\frac{(q-2)\left(\frac{p^*}{2}\right)'\left(\frac{p}{2\left(\frac{p^*}{2}\right)'}\right)'}{q}} \right. \\
 & \quad \left. \times |A_{k,t}^1|^{1 - \frac{(q-2)\left(\frac{p^*}{2}\right)'\left(\frac{p}{2\left(\frac{p^*}{2}\right)'}\right)'}{q}} \right]^{1 - \frac{2}{p}\left(\frac{p^*}{2}\right)'} \quad (65)
 \end{aligned}$$

Therefore, by (64) and (65), (63) becomes

$$\begin{aligned} \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |\tilde{A}|^2 \left(|(\text{adj}_2 Du)^1|^2 \right)^{\frac{q}{2}-1} dx &\leq 3 \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |A^1|^{p^*} dx \\ &+ 3 \left(\int_{B_R} |D\tilde{u}|^p dx \right)^{\frac{2}{p} \left(\frac{p^*}{2} \right)'} \left\{ \left(\int_{A_{k,t}^1} |(\text{adj}_2 Du)^1|^q dx \right)^{\frac{(q-2)}{q} \left(\frac{p^*}{2} \right)' \left(\frac{p}{2 \left(\frac{p^*}{2} \right)'} \right)'} \right. \\ &\quad \left. \times |A_{k,t}^1|^{1 - \frac{(q-2)}{q} \left(\frac{p^*}{2} \right)' \left(\frac{p}{2 \left(\frac{p^*}{2} \right)'} \right)'} \right\}^{1 - \frac{2}{p} \left(\frac{p^*}{2} \right)'}. \end{aligned} \quad (66)$$

Please, note that the previous condition $(q-2) \left(\frac{p^*}{2} \right)' \left(\frac{p}{2 \left(\frac{p^*}{2} \right)'} \right)' < q$ means $q < \frac{pp^*}{p+p^*}$. Finally, r -growth assumption (11) on $H(x, \cdot)$ yields

$$\int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu H(x, \det A) dx \leq \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu (k_3 |\det A|^r + c(x)) dx. \quad (67)$$

We compute $\det A$ with respect to the first row, see Lemma 3.6-(b),

$$\begin{aligned} \eta^\mu |\det A|^r &\leq \eta^\mu |A^1|^r |(\text{adj}_2 Du)^1|^r \leq (2\mu)^r \eta^{\mu-r} \left(\frac{u^1 - k}{t-s} \right)^r |(\text{adj}_2 Du)^1|^r \\ &\leq (2\mu)^r \left(\frac{u^1 - k}{t-s} \right)^r |(\text{adj}_2 Du)^1|^r. \end{aligned}$$

Notice that $r < p < p^*$ and $\frac{rp^*}{p^*-r} < q$. By the Young inequality with exponents $\frac{p^*}{r}$ and $\frac{p^*}{p^*-r}$, one has

$$\left(\frac{u^1 - k}{t-s} \right)^r |(\text{adj}_2 Du)^1|^r \leq \left(\frac{u^1 - k}{t-s} \right)^{p^*} + |(\text{adj}_2 Du)^1|^{\frac{rp^*}{p^*-r}}.$$

Hölder inequality with $\frac{q}{\frac{rp^*}{p^*-r}}$ and $\frac{q}{q-\frac{rp^*}{p^*-r}}$ leads to

$$\begin{aligned} \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu |\det A|^r dx &\leq (2\mu)^r \left[\int_{A_{k,t}^1 \cap \{\eta > 0\}} \left(\frac{u^1 - k}{t - s} \right)^{p^*} dx \right. \\ &\quad \left. + \int_{A_{k,t}^1 \cap \{\eta > 0\}} |(\operatorname{adj}_2 Du)^1|^{\frac{rp^*}{p^*-r}} dx \right] \\ &\leq (2\mu)^r \left[\int_{A_{k,t}^1 \cap \{\eta > 0\}} \left(\frac{u^1 - k}{t - s} \right)^{p^*} dx \right. \\ &\quad \left. + \left(\int_{B_R} |(\operatorname{adj}_2 Du)^1|^q dx \right)^{\frac{rp^*}{q(p^*-r)}} |A_{k,t}^1|^{1-\frac{rp^*}{q(p^*-r)}} \right]. \end{aligned} \quad (68)$$

Therefore, (67) and (68) imply

$$\begin{aligned} \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu H(x, \det A) dx &\leq k_3 (2\mu)^r \left[\int_{A_{k,t}^1 \cap \{\eta > 0\}} \left(\frac{u^1 - k}{t - s} \right)^{p^*} dx \right. \\ &\quad \left. + \left(\int_{B_R} |(\operatorname{adj}_2 Du)^1|^q dx \right)^{\frac{rp^*}{q(p^*-r)}} |A_{k,t}^1|^{1-\frac{rp^*}{q(p^*-r)}} \right] + \int_{A_{k,t}^1 \cap \{\eta > 0\}} \eta^\mu c(x) dx. \end{aligned} \quad (69)$$

By left-hand side inequalities in (9) and (11), using (58), (59), (60), (62), (66) and (69), we conclude

$$\begin{aligned} \int_{A_{k,t}^1} |Du^1|^p dx &\leq C \int_{A_{k,t}^1} \left(\frac{u^1 - k}{t - s} \right)^{p^*} dx + C \left\{ 1 + \|a + b + c\|_{L^\sigma(B_R)} \right. \\ &\quad \left. + \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\left(1-\frac{2}{p}\right)\left(\frac{p^*}{2}\right)'} \right. \\ &\quad \left. + \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{qp^*}{p(p^*-q)}} + \left(\int_{B_R} |(\operatorname{adj}_2 Du)^1|^q dx \right)^{\frac{rp^*}{q(p^*-r)}} \right. \\ &\quad \left. + 1_{(2,+\infty)}(q) \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{2}{p}\left(\frac{p^*}{2}\right)'} \right. \\ &\quad \left. \times \left(\int_{B_R} |(\operatorname{adj}_2 Du)^1|^q dx \right)^{\frac{(q-2)}{q}\left(\frac{p^*}{2}\right)'\left(\frac{p}{2\left(\frac{p^*}{2}\right)'}\right)\left(1-\frac{2}{p}\left(\frac{p^*}{2}\right)'\right)} \right\} |A_{k,t}^1|^\theta, \end{aligned} \quad (70)$$

where

$$\theta := \min \left\{ 1 - \left(1 - \frac{2}{p} \right) \left(\frac{p^*}{2} \right)', 1 - \frac{qp^*}{p(p^* - q)}, 1 - \frac{rp^*}{q(p^* - r)}, 1 - \frac{1}{\sigma}, \right. \\ \left. 1_{[1,2]}(q) + 1_{(2,+\infty)}(q) \left(1 - \frac{(q-2)}{q} \left(\frac{p^*}{2} \right)' \left(\frac{p}{2 \left(\frac{p^*}{2} \right)'} \right)' \right) \left(1 - \frac{2}{p} \left(\frac{p^*}{2} \right)' \right) \right\}$$

and $C = C(k_1, k_2, k_3, p, q, r, p^*) > 0$; moreover, $1_E(q) = 1$ if $q \in E$ and $1_E(q) = 0$ if $q \notin E$. Now we note that

$$1 - \left(1 - \frac{2}{p} \right) \left(\frac{p^*}{2} \right)' \geq 1 - \frac{p^*}{p(p^* - 1)} \geq 1 - \frac{qp^*}{p(p^* - q)},$$

where the last inequality is granted since $q \mapsto 1 - \frac{qp^*}{p(p^* - q)}$ decreases. Then,

$$\theta := \min \left\{ 1 - \frac{qp^*}{p(p^* - q)}, 1 - \frac{rp^*}{q(p^* - r)}, 1 - \frac{1}{\sigma}, \right. \\ \left. 1_{[1,2]}(q) + 1_{(2,+\infty)}(q) \left(1 - \frac{(q-2)}{q} \left(\frac{p^*}{2} \right)' \left(\frac{p}{2 \left(\frac{p^*}{2} \right)'} \right)' \right) \left(1 - \frac{2}{p} \left(\frac{p^*}{2} \right)' \right) \right\}.$$

Note that $\left(\frac{p^*}{2} \right)' = \frac{p^*}{p^* - 2}$; then, the exponents in (70) can be written as follows

$$\begin{aligned} \left(1 - \frac{2}{p} \right) \left(\frac{p^*}{2} \right)' &= \frac{p^*(p-2)}{p(p^* - 2)}, \quad \frac{2}{p} \left(\frac{p^*}{2} \right)' \\ &= \frac{2p^*}{p(p^* - 2)}, \quad \left(\frac{p^*}{2} \right)' \left(\frac{p}{2 \left(\frac{p^*}{2} \right)'} \right)' \left(1 - \frac{2}{p} \left(\frac{p^*}{2} \right)' \right) \\ &= \frac{p^*}{p^* - 2}, \end{aligned}$$

so that (70) turns out to be

$$\begin{aligned} \int_{A_{k,s}^1} |Du^1|^p dx &\leq C \int_{A_{k,t}^1} \left(\frac{u^1 - k}{t - s} \right)^{p^*} dx + C \left\{ 1 + \|a + b + c\|_{L^\sigma(B_R)} \right. \\ &\quad \left. + \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{p^*(p-2)}{p(p^* - 2)}} \right. \\ &\quad \left. + \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{qp^*}{p(p^* - q)}} \right\} \end{aligned}$$

$$\begin{aligned}
 & + \left(\int_{B_R} |(\operatorname{adj}_2 Du)^1|^q dx \right)^{\frac{rp^*}{q(p^*-r)}} \\
 & + 1_{(2,+\infty)}(q) \left(\int_{B_R} (|Du^2| + |Du^3|)^p dx \right)^{\frac{2p^*}{p(p^*-2)}} \\
 & \times \left(\int_{B_R} |(\operatorname{adj}_2 Du)^1|^q dx \right)^{\frac{(q-2)p^*}{q(p^*-2)}} \Big\} |A_{k,t}^1|^\theta, \quad (71)
 \end{aligned}$$

where

$$\begin{aligned}
 \theta &:= \min \left\{ 1 - \frac{qp^*}{p(p^*-q)}, 1 - \frac{rp^*}{q(p^*-r)}, 1 - \frac{1}{\sigma} \right\}, \quad \text{if } 1 < q \leq 2, \\
 \theta &:= \min \left\{ 1 - \frac{qp^*}{p(p^*-q)}, 1 - \frac{rp^*}{q(p^*-r)}, 1 - \frac{1}{\sigma}, \right. \\
 & \quad \left. \frac{p^*(p-2)-2p}{p(p^*-2)} - \frac{(q-2)p^*}{q(p^*-2)} \right\}, \quad \text{if } 2 < q.
 \end{aligned}$$

If $2 < q < \frac{pp^*}{p+p^*}$, we have

$$1 - \frac{qp^*}{p(p^*-q)} > \frac{p^*(p-2)-2p}{p(p^*-2)} - \frac{(q-2)p^*}{q(p^*-2)} = 1 - \frac{2p^*}{p(p^*-2)} - \frac{(q-2)p^*}{q(p^*-2)} \quad (72)$$

and

$$\theta = \min \left\{ 1 - \frac{2p^*}{p(p^*-2)} - \frac{(q-2)p^*}{q(p^*-2)}, 1 - \frac{rp^*}{q(p^*-r)}, 1 - \frac{1}{\sigma} \right\}.$$

This ends the proof of Proposition 4.1. \square

We now proceed with the proof of Theorem 2.2. We fix $x_0 \in \Omega$ and $R_0 < \min\{\operatorname{dist}(x_0, \partial\Omega), (\frac{3}{4\pi})^{1/3}\}$ such that

$$\int_{B_{R_0}} |u^1|^{p^*} dx < 1, \quad (73)$$

where B_ρ is the ball centered at x_0 with radius ρ . Note that $R_0 < 1$, $|B_{R_0}| < 1$ and $B_{R_0} \subset \subset \Omega$. For every $R \in (0, R_0]$ we define the decreasing sequence of radii

$$\rho_h := \frac{R}{2} + \frac{R}{2^{h+1}}.$$

Fix a positive constant $d \geq 1$ and define the increasing sequence of positive levels

$$k_h := d \left(1 - \frac{1}{2^{h+1}} \right).$$

We define the “excess”

$$J_h := \int_{A_{k_h, \rho_h}^1} (u^1 - k_h)^{p^*} dx.$$

We use our Caccioppoli inequality (48) and Proposition 2.4 of [13]: we get

$$J_{h+1} \leq c \left(2^{\frac{p^* p^*}{p}} \right)^h (J_h)^{\theta \frac{p^*}{p}},$$

where the positive constant c is independent of h . See also [40, 41]. Assumption (17) tells us that $\theta \frac{p^*}{p} > 1$; then, we can use Lemma 2.5 of [13] with $\gamma := \theta \frac{p^*}{p} - 1$, see also [42]:

$$J_h \leq \left(2^{\frac{p^* p^*}{p}} \right)^{-\frac{h}{\gamma}} J_0, \quad (74)$$

provided

$$J_0 \leq c^{-\frac{1}{\gamma}} \left(2^{\frac{p^* p^*}{p}} \right)^{-\frac{1}{\gamma^2}}. \quad (75)$$

Note that

$$J_0 = \int_{A_{\frac{d}{2}, R}^1} \left(u^1 - \frac{d}{2} \right)^{p^*} dx \rightarrow 0 \quad \text{as} \quad d \rightarrow +\infty;$$

then, we can choose $d \geq 1$ large enough so that (75) holds true. Thus, we have (74) with $\gamma > 0$, so that $J_h \rightarrow 0$ as $h \rightarrow +\infty$; since $\frac{R}{2} < \rho_h$ and $k_h < d$, we also have

$$0 \leq \int_{A_{d, \frac{R}{2}}^1} (u^1 - d)^{p^*} dx \leq J_h;$$

then,

$$\int_{A_{d, \frac{R}{2}}^1} (u^1 - d)^{p^*} dx = 0$$

so that $u^1 \leq d$ almost everywhere in $B_{\frac{R}{2}}$. We have proved that u^1 is locally bounded from above. In order to prove that u^1 is locally bounded from below, we note that $-u$ locally minimizes $\int_{\Omega} \tilde{f}(x, Dz(x)) dx$, where $\tilde{f}(x, \xi) = f(x, -\xi)$; then, we get that $-u^1$ is locally bounded from above, so u^1 is locally bounded from below. We have just shown that $u^1 \in L_{\text{loc}}^{\infty}(\Omega)$.

Now we turn our attention to the second component u^2 . We change the order of the two components u^1 and u^2 : we get a new function v as follows:

$$v = \begin{pmatrix} u^2 \\ u^1 \\ u^3 \end{pmatrix};$$

then,

$$Dv = \begin{pmatrix} Du^2 \\ Du^1 \\ Du^3 \end{pmatrix}$$

and $\det Dv = -\det Du$; moreover $(\operatorname{adj}_2 Dv)^1 = -(\operatorname{adj}_2 Du)^2$, $(\operatorname{adj}_2 Dv)^2 = -(\operatorname{adj}_2 Du)^1$ and $(\operatorname{adj}_2 Dv)^3 = -(\operatorname{adj}_2 Du)^3$, so that

$$\operatorname{adj}_2 Dv = - \begin{pmatrix} (\operatorname{adj}_2 Du)^2 \\ (\operatorname{adj}_2 Du)^1 \\ (\operatorname{adj}_2 Du)^3 \end{pmatrix}.$$

If we write $C_{1,2}(\xi)$ to denote the matrix obtained from ξ by inverting line 1 and line 2, we have $Dv = C_{1,2}(Du)$ and $\operatorname{adj}_2 Dv = -C_{1,2}(\operatorname{adj}_2 Du)$. Then, v is a local minimizer of $\int_{\Omega} \tilde{f}(x, Dw(x))dx$, where $\tilde{f}(x, \xi) = f(x, C_{1,2}(\xi))$. Thus, the first component v^1 is locally bounded: $u^2 = v^1 \in L_{\text{loc}}^{\infty}(\Omega)$. In a similar way we deal with the third component u^3 : we change the order of the two components u^1 and u^3 ; we get a new function w as follows:

$$w = \begin{pmatrix} u^3 \\ u^2 \\ u^1 \end{pmatrix};$$

then,

$$Dw = \begin{pmatrix} Du^3 \\ Du^2 \\ Du^1 \end{pmatrix}$$

and $\det Dw = -\det Du$; moreover $(\operatorname{adj}_2 Dw)^1 = -(\operatorname{adj}_2 Du)^3$, $(\operatorname{adj}_2 Dw)^2 = -(\operatorname{adj}_2 Du)^2$ and $(\operatorname{adj}_2 Dw)^3 = -(\operatorname{adj}_2 Du)^1$, so that

$$\operatorname{adj}_2 Dw = - \begin{pmatrix} (\operatorname{adj}_2 Du)^3 \\ (\operatorname{adj}_2 Du)^2 \\ (\operatorname{adj}_2 Du)^1 \end{pmatrix}.$$

If we write $C_{1,3}(\xi)$ to denote the matrix obtained from ξ by inverting line 1 and line 3, we have $Dw = C_{1,3}(Du)$ and $\operatorname{adj}_2 Dw = -C_{1,3}(\operatorname{adj}_2 Du)$. Then, w is a

local minimizer of $\int_{\Omega} \tilde{\tilde{f}}(x, Dz(x))dx$, where $\tilde{\tilde{f}}(x, \xi) = f(x, C_{1,3}(\xi))$. Thus, the first component w^1 is locally bounded: $u^3 = w^1 \in L_{\text{loc}}^{\infty}(\Omega)$. This ends the proof of Theorem 2.2. \square

5 Conclusions

We have been able to prove boundedness for minimizers of the most important three-dimensional polyconvex integral, provided the growth exponents verify some restrictions. It would be interesting to understand what happens when such restrictions are not in force.

Acknowledgements We thank the referee for carefully reading the manuscript and for the useful remarks. M. Carozza, R. Gioia and F. Leonetti have been supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). H. Gao thanks NSFC (10371050) and NSF of Hebei Province (A2015201149) for their support. R. Gioia has been partially supported by Università degli Studi di Napoli "Parthenope" through the Project "Sostegno alla ricerca individuale (annualità 2015-2016-2017)" and the Project "Sostenibilità, eternalità e uso efficiente delle risorse ambientali"(triennio 2017-2019). F. Leonetti acknowledges also the support of UNIVAQ.

Appendix: Comparison Between Two Structures

Lemma A.1 *We assume that $F^{\alpha}, G^{\alpha} : \mathbb{R}^3 \mapsto [0, +\infty[$ and $H : \mathbb{R} \mapsto [0, +\infty[$; let $p, q, r \in]0, +\infty[$ with $p \neq 2$. Then, it is false that*

$$\sum_{\alpha=1}^3 F^{\alpha}(\xi^{\alpha}) + \sum_{\alpha=1}^3 G^{\alpha}((\text{adj}_2 \xi)^{\alpha}) + H(\det \xi) = |\xi|^p + |\text{adj}_2 \xi|^q + |\det \xi|^r \quad (76)$$

for every $\xi \in \mathbb{R}^{3 \times 3}$.

Proof We argue by contradiction: if (76) holds true, then we can use (76) with

$$\xi = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (77)$$

and we get

$$\text{adj}_2 \xi = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (78)$$

with $\det \xi = 0$, so that

$$\sum_{\alpha=1}^3 F^{\alpha}((0, 0, 0)) + \sum_{\alpha=1}^3 G^{\alpha}((0, 0, 0)) + H(0) = 0; \quad (79)$$

we keep in mind that $F^\alpha, G^\alpha, H \geq 0$ and we get

$$F^\alpha((0, 0, 0)) = G^\alpha((0, 0, 0)) = H(0) = 0, \quad (80)$$

for every $\alpha = 1, 2, 3$. Now we use (76) with

$$\xi = \begin{pmatrix} t & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (81)$$

and we get

$$\text{adj}_2 \xi = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (82)$$

with $\det \xi = 0$, so that

$$F^1((t, 0, 0)) + F^2((0, 0, 0)) + F^3((0, 0, 0)) + \sum_{\alpha=1}^3 G^\alpha((0, 0, 0)) + H(0) = |t|^p; \quad (83)$$

we keep in mind (80) and we get

$$F^1((t, 0, 0)) = |t|^p, \quad (84)$$

for every $t \in \mathbb{R}$. In a similar manner, taking

$$\xi = \begin{pmatrix} 0 & 0 & 0 \\ t & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (85)$$

we get

$$F^2((t, 0, 0)) = |t|^p, \quad (86)$$

for every $t \in \mathbb{R}$. In the same way, taking

$$\xi = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ t & 0 & 0 \end{pmatrix}, \quad (87)$$

we get

$$F^3((t, 0, 0)) = |t|^p, \quad (88)$$

for every $t \in \mathbb{R}$. Eventually, we take

$$\xi = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad (89)$$

and (76) implies

$$\sum_{\alpha=1}^3 F^{\alpha}((1, 0, 0)) + \sum_{\alpha=1}^3 G^{\alpha}((0, 0, 0)) + H(0) = 3^{p/2}; \quad (90)$$

we use (80), (84), (86), (88) and we get

$$3 = 3^{p/2} : \quad (91)$$

such an equality is a contradiction, since $p \neq 2$. This ends the proof of Lemma A.1. \square

References

1. Mingione, G.: Regularity of minima: an invitation to the dark side of the calculus of variations. *Appl. Math.* **51**, 355–426 (2006)
2. Mingione, G.: Singularities of minima: a walk in the wild side of the calculus of variations. *J. Glob. Optim.* **40**, 209–223 (2008)
3. Fusco, N., Hutchinson, J.: Partial regularity in problems motivated by nonlinear elasticity. *SIAM J. Math. Anal.* **22**, 1516–1551 (1991)
4. Fuchs, M., Seregin, G.: Partial regularity of the deformation gradient for some model problems in nonlinear twodimensional elasticity. *Algebra i Analiz* **6**, 128–153 (1994)
5. Fuchs, M., Reuling, J.: Partial regularity for certain classes of polyconvex functionals related to non linear elasticity. *Manusc. Math.* **87**, 13–26 (1995)
6. Passarelli di Napoli, A.: A regularity result for a class of polyconvex functionals. *Ricerche Mat.* **48**, 379–393 (1999)
7. Esposito, L., Mingione, G.: Partial regularity for minimizers of degenerate polyconvex energies. *J. Convex Anal.* **8**, 1–38 (2001)
8. Hamburger, C.: Partial regularity of minimizers of polyconvex variational integrals. *Calc. Var. Partial Differ. Equ.* **18**, 221–241 (2003)
9. Carozza, M., Passarelli di Napoli, A.: Model problems from nonlinear elasticity: partial regularity results. *ESAIM Control Opt. Calc. Var.* **13**, 120–134 (2007)
10. Carozza, M., Leone, C., Passarelli di Napoli, A., Verde, A.: Partial regularity for polyconvex functionals depending on the Hessian determinant. *Calc. Var. Partial Differ. Equ.* **35**, 215–238 (2009)
11. Fusco, N., Hutchinson, J.: Partial regularity and everywhere continuity for a model problem from non-linear elasticity. *J. Aust. Math. Soc. Ser. A* **57**, 158–169 (1994)
12. Fuchs, M., Seregin, G.: Holder continuity for weak estremals of some two-dimensional variational problems related to non linear elasticity. *Adv. Math. Sci. Appl.* **7**, 413–425 (1997)
13. Cupini, G., Leonetti, F., Mascolo, E.: Local boundedness for minimizers of some polyconvex integrals. *Arch. Ration. Mech. Anal.* **224**, 269–289 (2017)
14. Leonetti, F.: Maximum principle for vector-valued minimizers of some integral functionals. *Boll. Unione Mat. Ital. A* **5**, 51–56 (1991)
15. D’Ottavio, A., Leonetti, F., Musciano, C.: Maximum principle for vector-valued mappings minimizing variational integrals. *Atti Sem. Mat. Fis. Modena* **46 Suppl.**, 677–683 (1998)
16. Leonetti, F., Siepe, F.: Maximum principle for vector-valued minimizers of some integral functionals. *J. Convex Anal.* **12**, 267–278 (2005)
17. Leonetti, F., Siepe, F.: Bounds for vector valued minimizers of some integral functionals. *Ricerche Mat.* **54**, 303–312 (2005)
18. Leonetti, F., Petricca, P.V.: Bounds for some minimizing sequences of functionals. *Adv. Calc. Var.* **4**, 83–100 (2011)
19. Leonetti, F., Petricca, P.V.: Bounds for vector valued minimizers of some relaxed functionals. *Complex Var. Elliptic Equ.* **58**, 221–230 (2013)

20. Sverák, V.: Regularity properties for deformations with finite energy. *Arch. Ration. Mech. Anal.* **100**, 105–127 (1988)
21. Bauman, P., Owen, N., Phillips, D.: Maximum principles and a priori estimates for a class of problems from nonlinear elasticity. *Ann. Inst. Henri Poincaré Anal. Non Linéaire* **8**, 119–157 (1991)
22. Bauman, P., Phillips, D., Owen, N.: Maximal smoothness of solutions to certain Euler–Lagrange equations from nonlinear elasticity. *Proc. R. Soc. Edinb. Sect. A* **119**, 241–263 (1991)
23. Bauman, P., Owen, N., Phillips, D.: Maximum principles and a priori estimates for an incompressible material in nonlinear elasticity. *Commun. Partial Differ. Equ.* **17**, 1185–1212 (1992)
24. Bauman, P., Phillips, D.: Univalent minimizers of polyconvex functionals in two dimensions. *Arch. Ration. Mech. Anal.* **126**, 161–181 (1994)
25. Bevan, J.: A condition for the Hölder regularity of local minimizers of a nonlinear elastic energy in two dimensions. *Arch. Ration. Mech. Anal.* **225**, 249–285 (2017)
26. Leonetti, F.: Pointwise estimates for a model problem in nonlinear elasticity. *Forum Math.* **18**, 529–534 (2006)
27. Gao, H., Leonetti, F., Macri, M., Petricca, P.V.: Regularity for minimizers with positive Jacobian. Preprint (2017)
28. Ball, J.: Convexity conditions and existence theorems in nonlinear elasticity. *Arch. Ration. Mech. Anal.* **63**, 337–403 (1977)
29. Dacorogna, B.: *Direct Methods in the Calculus of Variations*, 2nd edn. Appl. Math. Sci. vol. 78. Springer, New York (2008)
30. De Giorgi, E.: An example of a discontinuous solution to an elliptic variational problem. *Boll. Unione Mat. Ital.* **4**(1), 135–137 (1968). (in Italian)
31. Sverák, V., Yan, X.: A singular minimizer of a smooth strongly convex functional in three dimensions. *Calc. Var. Partial Differ. Equ.* **10**, 213–221 (2000)
32. Mooney, C., Savin, O.: Some singular minimizers in low dimensions in the calculus of variations. *Arch. Ration. Mech. Anal.* **221**, 1–22 (2016)
33. Giaquinta, M.: Growth conditions and regularity, a counterexample. *Manuscr. Math.* **59**, 245–248 (1987)
34. Marcellini, P.: Un exemple de solution discontinue d'un problème variationnel dans le cas scalaire. Preprint 11, Istituto Matematico “U. Dini” Università di Firenze (1987)
35. Marcellini, P.: Regularity of minimizers of integrals of the calculus of variations with non standard growth conditions. *Arch. Ration. Mech. Anal.* **105**, 267–284 (1989)
36. Marcellini, P.: Regularity and existence of solutions of elliptic equations with $p - q$ -growth conditions. *J. Differ. Equ.* **90**, 1–30 (1991)
37. Hong, M.C.: Some remarks on the minimizers of variational integrals with non standard growth conditions. *Boll. Un. Mat. Ital. A* **6**, 91–101 (1992)
38. Esposito, L., Leonetti, F., Mingione, G.: Sharp regularity results for functionals with (p, q) growth. *J. Differ. Equ.* **204**, 5–55 (2004)
39. Fonseca, I., Maly, J., Mingione, G.: Scalar minimizers with fractal singular sets. *Arch. Ration. Mech. Anal.* **172**, 295–307 (2004)
40. Moscarillo, G., Nania, L.: Hölder continuity of minimizers of functionals with non standard growth conditions. *Ricerche Mat.* **40**, 259–273 (1991)
41. Fusco, N., Sbordone, C.: Some remarks on the regularity of minima of anisotropic integrals. *Commun. Partial Differ. Equ.* **18**, 153–167 (1993)
42. Giusti, E.: *Direct Methods in the Calculus of Variations*. World Scientific Publishing, Singapore (2003)